



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

in the number of the staminal whorls. Calyx and corolla present the normal condition, while the stamens have taken the shape of petals. Though the ten stamens belong to two whorls, the petalodes are more or less connate at the base into 10-merous whorls; but the petalodes of all the whorls are in superposition, showing thus that they must be considered as consisting of alternate 5-merous whorls, and this is also apparent by the aestivation of the broader petalodes of the outer whorls; the first five petalodes, of course, are opposite to the corolla lobes, as the ericaceous flower is obdiplostemonous. The number of 10-merous whorls in each flower is about ten; the shape of the petalodes changes from the broadly oval of the outer ones to the oblong shape of the innermost petalodes. Even the carpels are changed into narrow oblong petalodes. Only in one case I found an incompletely developed anther adhering to one of the petalodes.

As it would be interesting to have this shrub in cultivation, though the horticultural value of this form is lessened by the rather difficult cultivation of the species, we sent rooted suckers to the Arnold Arboretum, but the plants did not survive, and concerning the fate of a few sent to a German nursery I know nothing. The old plant, however, is in all probability still there, and we may yet hope to see this handsome form introduced into cultivation. Dried specimens of it are preserved in the herbarium of the Arnold Arboretum.—ALFRED REHDER, *Arnold Arboretum*.

PUCCINIA UPON MELOTHRIA

(WITH ONE FIGURE)


This rust was found in the neighborhood of West Raleigh, N. C., in the autumn of 1907 upon *Melothria pendula*. The host is not a common plant in this locality, only two plants being seen during an entire collecting season; one of these was thoroughly infected, the other not at all.

The species is of special interest on account of the comparative paucity of rusts affecting the cucurbits, as well as owing to the economic importance of this family. On the whole family as represented by the North American genera, either native or exotic, listed in SMALL'S *Flora of the South Eastern United States*, GRAY'S *Manual*, GRAY'S *Field, Forest, and Garden Botany*, and BRITTON and BROWN'S *Illustrated Flora*, there are mentioned in FARLOW'S *Host Index* no rusts at all. In SACCARDO there are six species, as follows: (1) *Uromyces Melothriae* on *M. tomentosa* in Abyssinia; (2) *U. Cayaponiae* on *C. racemosa* in Africa; (3) *U. Hellarianus* on *Cayaponia* in Porto Rico; (4) *Puccinia Cucumeris* on *C. ficifolia* in Abyssinia; (5) *P.*

Momordicæ on *M. cordifolia* in Natal; (6) *Aecidium Momordicæ* on *Momordica* in Brazil.

The *Puccinia* upon *Melothria pendula* is different in shape, color, and size of spores from *P. Cucumeris* and from *P. Momordicæ*, and seems clearly to be an undescribed species. The germination of the teleuto-spores occurs freely in the older sori, placing the rust among the *Leptopuccinia*. The following description is proposed.

***Puccinia Melothriae*, n. sp. — III.** Teleutosori: hypophyllous, crowded in irregular concentric circles, pustular, soon naked, pulverulent, cinnamon-colored (no. 32 Sacc.), cinereous on germination, ruptured epidermis inconspicuous; spots pale, indefinite. Teleutospores: hyaline to straw-colored (no. 26 Sacc.), $40-52 \times 10-16 \mu$; slightly or not at all thickened at apex, often not constricted at septum, wall about 2μ thick, smooth. Terminal cell usually round, rarely oblong, $10-16 \times 12-20 \mu$. Basal cell wedge-shaped, $10-15 \times 20-30 \mu$. Pedicel $20-45 \mu$ long, hyaline. Germination immediate; that of the terminal cell by an apical promycelium, that of the basal cell by a lateral promycelium.



Puccinia Melothriae, n. sp.

On *Melothria pendula* L., West Raleigh, N. C., 9-15-06, F. L. Stevens and J. G. Hall, type no. 471.—F. L. STEVENS, West Raleigh, N. C.

AN INSTANCE OF MUTATION

Euphorbia marginata Pursh, commonly called in Colorado "snow-on-the-mountain," is easily recognized by its white-margined upper leaves or bracts, and the five white appendages of the floral glands. In the splitting up of the old genus *Euphorbia*, it has been referred to a genus *Dichrophyllum* Kl. & Garcke, and distinguished from *Tithymalus* Adans by the characters just mentioned. In *Tithymalus* there are but four glands, with rudiments of the fifth. All authors seem to agree that *E. marginata* has five glands; "three" in BRITTON and BROWN, *Ill. Flora* 2:376, is a mere misprint, as the figure shows. Having recently had occasion to examine *E. marginata* at Boulder, Colorado, I have been astonished to find that *four* glands is normal in this locality, and a plant with *five* glands